

Overview

The Contribution of Cytotoxic Chemotherapy to 5-year Survival in Adult Malignancies

Graeme Morgan*, Robyn Ward†, Michael Barton‡

Table 2 — Impact of cytotoxic chemotherapy on 5-year survival in American adults

Malignancy	ICD-9	Number of cancers in people aged > 20 years*	Absolute number of 5-year survivors due to chemotherapy†	Percentage 5-year survivors due to chemotherapy‡
Head and neck	140-149, 160, 161	5139	97	1.9
Esophagus	150	1521	82	4.9
Stomach	151	3001	20	0.7
Colon	153	13 936	146	1.0
Rectum	154	5533	189	3.4
Pancreas	157	3567	—	—
Lung	162	20 741	410	2.0
Soft tissue sarcoma	171	858	—	—
Melanoma	172	8546	—	—
Breast	174	31 133	446	1.4
Uterus	179-182	4611	—	—
Cervix	180	1825	219	12
Ovary	183	3032	269	8.9
Prostate	185	23 242	—	—
Testis	186	989	373	37.7
Bladder	188	6667	—	—
Kidney	189	3722	—	—
Brain	191	1824	68	3.7
Unknown primary site	195-199	6200	—	—
Non-Hodgkin's lymphoma	200 + 202	6217	653	10.5
Hodgkin's disease	201	846	341	40.3
Multiple myeloma	203	1721	—	—
Total		154 971	3306	2.1%

*Numbers from Ref. [22].

†Absolute numbers (see text).

‡% for individual malignancy.